

VOLUME OF REPRESENTATION VARIETIES

MOTOHICO MULASE¹ AND MICHAEL PENKAVA²

ABSTRACT. We introduce the notion of volume of the representation variety of a finitely presented discrete group in a compact Lie group using the push-forward measure associated to a map defined by a presentation of the discrete group. We show that the volume thus defined is invariant under the Andrews-Curtis moves of the generators and relators of the discrete group, and moreover, that it is actually independent of the choice of presentation if the difference of the number of generators and the number of relators remains the same. We then calculate the volume of the representation variety of a surface group in an arbitrary compact Lie group using the classical technique of Frobenius and Schur on finite groups. Our formulas recover the results of Witten and Liu on the symplectic volume and the Reidemeister torsion of the moduli space of flat G -connections on a surface up to a constant factor when the Lie group G is semisimple.

CONTENTS

0. Introduction and the Main Results	1
1. Presentation of a Group	4
2. Volume of Representation Varieties	6
3. Representation Varieties of Orientable Surface Groups	13
4. Representation Varieties of Non-orientable Surface Groups	16
5. Harmonic Analysis and Irreducible Representations of a Compact Lie Group	20
6. Representation Varieties and Cell-Decompositions of a Surface	23
References	26

0. INTRODUCTION AND THE MAIN RESULTS

Let Π be a finitely presented discrete group generated by k elements with r relations:

$$(0.1) \quad \Pi = \langle a_1, \dots, a_k \mid q_1, \dots, q_r \rangle .$$

The space $\text{Hom}(\Pi, G)$ of homomorphisms from Π into a compact Lie group G is called the *representation variety* of Π in G . Representation varieties of the fundamental group of a manifold appear in various places in low-dimensional geometry and topology (see for example, [8, 14, 23, 24, 43]). To the presentation (0.1) we associate a *presentation map*

$$(0.2) \quad q : G^k \ni x = (x_1, \dots, x_k) \longmapsto (q_1(x), \dots, q_r(x)) \in G^r ,$$

which is a real analytic map. By definition, we have a canonical identification

$$(0.3) \quad \text{Hom}(\Pi, G) = q^{-1}(1, \dots, 1)$$

Date: December 1, 2002.

¹Research supported by NSF grant DMS-9971371 and the University of California, Davis.

²Research supported by NSF grant DMS-0200669 and the University of Wisconsin, Eau Claire.

that gives a realization of the representation variety as a real analytic subvariety of G^k . If the relators are given without redundancy, then its dimension is expected to be

$$(0.4) \quad \dim \operatorname{Hom}(\Pi, G) = (k - r) \dim G .$$

The purpose of this paper is to give a definition of the *volume* $|\operatorname{Hom}(\Pi, G)|$ of the representation variety and to study some of its properties. Since we use the push-forward measure via the presentation map (0.2) to define the volume, the central question is to determine whether it is an invariant of the group. In this article we give an affirmative answer to this question. The main results of the paper are the following.

Theorem 0.1. *The volume is invariant under the Andrews-Curtis moves.*

Because of the dimension formula of (0.4), it is necessary that the difference $k - r$ of the numbers of generators and relators remains invariant for different presentations of the group Π to give the same volume $|\operatorname{Hom}(\Pi, G)|$. Surprisingly, the condition is sufficient.

Theorem 0.2. *The volume is independent of the choice of presentation if the difference between the number of generators and the number of relators is the same.*

The volume of the representation variety of the fundamental group of a closed surface is a topological invariant and actually expressible in terms of irreducible representations of the compact Lie group G .

Theorem 0.3. *Let S be a closed orientable surface of genus $g \geq 2$. Then the volume of the representation variety can be computed by the formula*

$$(0.5) \quad \sum_{\lambda \in \hat{G}} (\dim \lambda)^{\chi(S)} = |G|^{\chi(S)-1} |\operatorname{Hom}(\pi_1(S), G)|$$

if the sum of the LHS converges, where $\hat{G}$ denotes the set of isomorphism classes of complex irreducible representations of G , $\dim \lambda$ is the complex dimension of the irreducible representation λ , and $\chi(S) = 2 - 2g$ is the Euler characteristic of S . Since G is compact, $\hat{G}$ is a countable set.

For a non-orientable surface S , the formula for the volume involves more detailed information on the irreducible representations of G . Using the Frobenius-Schur indicator of irreducible characters [12], we decompose the set of complex irreducible representations $\hat{G}$ into the union of three disjoint subsets, corresponding to real, complex, and quaternionic irreducible representations:

$$(0.6) \quad \begin{aligned} \hat{G}_1 &= \left\{ \lambda \in \hat{G} \mid \frac{1}{|G|} \int_G \chi_\lambda(w^2) dw = 1 \right\} ; \\ \hat{G}_2 &= \left\{ \lambda \in \hat{G} \mid \frac{1}{|G|} \int_G \chi_\lambda(w^2) dw = 0 \right\} ; \\ \hat{G}_4 &= \left\{ \lambda \in \hat{G} \mid \frac{1}{|G|} \int_G \chi_\lambda(w^2) dw = -1 \right\} , \end{aligned}$$

where χ_λ is the irreducible character of $\lambda \in \hat{G}$. Now we have

Theorem 0.4. *Let S a closed non-orientable surface. Then we have*

$$(0.7) \quad \sum_{\lambda \in \hat{G}_1} (\dim \lambda)^{\chi(S)} + \sum_{\lambda \in \hat{G}_4} (-\dim \lambda)^{\chi(S)} = |G|^{\chi(S)-1} |\operatorname{Hom}(\pi_1(S), G)|$$

if the sum of the LHS is absolutely convergent.

The formulas (0.5) and (0.7) for a finite group, in which case the closed surface can be anything, are due to Mednykh [29] and Frobenius-Schur [12], respectively. We refer to the excellent review article [21] for the history of related topics. For a finite group G and $S = S^2$, (0.5) reduces to the classical formula for the order of the group

$$(0.8) \quad |G| = \sum_{\lambda \in \hat{G}} (\dim \lambda)^2 .$$

For a compact connected semisimple Lie group G , the formulas agree with the results of Witten [52, 54] and Liu [25, 26, 27] up to a constant factor. They calculated the natural symplectic volume and the Reidemeister torsion of the moduli space of flat G -connections on a surface S using techniques from theoretical physics and analysis of heat kernels. The idea of using Chern-Simons gauge theory to obtain (0.5) for a finite group was carried out in [10]. The representation variety has algebro-geometric interest as the moduli space of holomorphic vector bundles on a compact Riemann surface, and through this interpretation it has deep connections with conformal field theory and topological quantum field theory. Let S be a compact Riemann surface with a complex structure. Narasimhan-Seshadri theory [37, 38] shows that the moduli space of stable holomorphic $G_{\mathbb{C}}$ bundles over S as a real analytic variety can be identified with

$$\mathcal{M}(S, G_{\mathbb{C}}) = \frac{\text{Hom}(\pi_1(S), G)}{G/Z(G)} ,$$

where $Z(G)$ is the center of G and the G -action on $\text{Hom}(\pi_1(S), G)$ is through conjugation. The first Chern class of the determinant line bundle over the moduli space gives an integral symplectic form ω on the moduli space. Using this 2-form Witten [52] computed the natural symplectic volume of $\mathcal{M}(S, G_{\mathbb{C}})$. His formula and ours differ by a factor of $(2\pi)^{\dim_{\mathbb{R}} \mathcal{M}}$, which is due to the normalization of the integral cohomology class $[\omega]$ on the moduli space, but the factor does not come from the choice of the Haar measure on the group G . From the algebro-geometric point of view, certainly the quotient of the natural conjugate action of the representation variety is a desirable object of study. However, our formulas (0.5) and (0.7) suggest that the volume of the whole representation variety measured in comparison with the volume of the space $G^{1-\chi(S)}$ may be a more natural object. The heat kernel method is necessary to connect the volume of the representation variety of a surface group we discuss in this paper to the symplectic volume or the Reidemeister torsion of the moduli space of flat connections on a closed surface. Our point is that if we consider the natural quantity $|G|^{\chi(S)-1} |\text{Hom}(\pi_1(S), G)|$, which is independent of the choice of the measure on G , then we can compute its value by exactly the same classical method of Frobenius [11] and Frobenius-Schur [12] for finite groups.

The paper is organized as follows. In Section 1 we recall that a group Π is completely determined by the hom functor $\text{Hom}(\Pi, \bullet)$. This implies that two presentations define the same group if and only if what we call the presentation functors are naturally isomorphic. With this preparation we define the volume of the representation variety $\text{Hom}(\Pi, G)$ in Section 2 using a presentation of Π . We then prove that the volume is actually independent of the presentation, utilizing the behavior of the volume under the Andrews-Curtis moves. Then in Sections 3 and 4 we calculate the volume of the representation variety of a surface group, the orientable case first followed by the non-orientable case. Throughout the paper except for Section 1 we need non-commutative harmonic analysis of distributions on a

compact Lie group and character theory. These techniques are summarized in Section 5. The final section is devoted to presenting a heuristic argument that relates the computation of the volume of this paper to the non-commutative matrix integral studied in [35, 36].

Acknowledgement. The authors thank Bill Goldman for drawing their attention to important references on the subject of this article. They also thank Greg Kuperberg, Abby Thompson and Bill Thurston for stimulating discussions on representation varieties and Andrews-Curtis moves.

1. PRESENTATION OF A GROUP

Consider a set of *alphabets* $\{a_1, \dots, a_k\}$ consisting of k letters. A *word* $W(a)$ of length ν is the expression of the form

$$(1.1) \quad W(a) = a_{w(1)}^{\epsilon(1)} \cdots a_{w(\nu)}^{\epsilon(\nu)},$$

where the subscript

$$(1.2) \quad w : \{1, \dots, \nu\} \longrightarrow \{1, \dots, k\}$$

indicates the ℓ -th letter $a_{w(\ell)}$ in the word $W(a)$, and the exponent $\epsilon(\ell) = \pm 1$ represents either the letter $a_{w(\ell)}$ or its *inverse* $a_{w(\ell)}^{-1}$. Instead of the commonly used expression a_i^ν , we always use

$$\overbrace{a_i \cdots a_i}^\nu,$$

which is a word of length ν . Throughout this article, every word is assumed to be reduced, namely, no expressions like $a_i a_i^{-1}$ and $a_i^{-1} a_i$ appear in a word.

For every given group G , a word $W(a)$ of letters $a_1, \dots, a_k$ (that are unrelated with G) defines a map

$$(1.3) \quad W : G^k \ni x = (x_1, \dots, x_k) \longmapsto W(x) = x_{w(1)}^{\epsilon(1)} \cdots x_{w(\nu)}^{\epsilon(\nu)} \in G.$$

The map W is the composition of two maps

$$(1.4) \quad \begin{array}{ccc} G^k & \ni & (x_1, \dots, x_k) \\ \downarrow \iota_W & & \downarrow \iota_W \\ G^\nu & \ni & (x_{w(1)}^{\epsilon(1)}, \dots, x_{w(\nu)}^{\epsilon(\nu)}) \\ \downarrow m & & \downarrow m \\ G & \ni & x_{w(1)}^{\epsilon(1)} \cdots x_{w(\nu)}^{\epsilon(\nu)}, \end{array}$$

where $\iota_W : G^k \rightarrow G^\nu$ represents the shape of the word $W(a)$, and

$$(1.5) \quad m : G^\nu \ni (y_1, \dots, y_\nu) \longmapsto y_1 \cdots y_\nu \in G$$

is the multiplication map. If the word $W(a)$ does not contain the letter a_i , then x_i is sent to $1 \in G$ by the map ι_W . In particular, if $W = \emptyset$ is the empty word, then $\iota_\emptyset$ is the trivial map that sends everything in G^k to $\{1\} = G^0$.

Let us consider a finitely presented discrete group

$$(1.6) \quad \Pi = \langle a_1, \dots, a_k \mid q_1, \dots, q_r \rangle$$

defined by k generators and r relations $q_1 = \cdots = q_r = 1$. Every *relator* q_j is a reduced word, and we always assume that none of the relators is the empty word. The set of relators defines the *presentation map*

$$(1.7) \quad q : G^k \ni x = (x_1, \dots, x_k) \longmapsto (q_1(x), \dots, q_r(x)) \in G^r .$$

We give a group structure to the target space G^r as the product group of r copies of G . The identity element of G^r is $1 = (1, \dots, 1)$. The *representation space* of Π in G is the set of group homomorphisms from Π to G and is naturally identified with

$$(1.8) \quad \text{Hom}(\Pi, G) = q^{-1}(1) .$$

Let $\mathcal{G}$ denote the category of groups with group homomorphisms as morphisms among the objects. A finite presentation (1.6) defines a covariant functor from $\mathcal{G}$ to the category $\mathcal{S}$ of sets, associating $q^{-1}(1)$ to a group G . Let us call this functor the *presentation functor*. The covariance of the presentation functor is a consequence of the commutativity of the diagram

$$(1.9) \quad \begin{array}{ccc} H^k & \xrightarrow{q} & H^r \\ \gamma \downarrow & & \downarrow \gamma \\ G^k & \xrightarrow{q} & G^r \end{array}$$

for every homomorphism $\gamma : H \rightarrow G$, where the vertical arrows are the natural maps induced by γ . This functor is representable and the group Π is the representing object in $\mathcal{G}$ [28]. Thus the presentation functor, whose representation is the hom functor $\text{Hom}(\Pi, \bullet)$, is determined by the group Π alone and is independent of the choice of its presentation. Let $\mu : \Pi \rightarrow \Pi'$ be a group homomorphism from Π to Π' . The pull-back map μ^* defines the natural morphism

$$\mu^* : \text{Hom}(\Pi', \bullet) \longrightarrow \text{Hom}(\Pi, \bullet)$$

between the hom functors.

Notice that the functor from $\mathcal{G}$ to the functor category $\mathcal{F}(\mathcal{G}, \mathcal{S})$ that associates

$$\mathcal{G} \ni \Pi \longmapsto \text{Hom}(\Pi, \bullet) \in \mathcal{F}(\mathcal{G}, \mathcal{S})$$

is a faithful contravariant functor. Indeed, if there is a natural isomorphism

$$A : \text{Hom}(\Pi, \bullet) \xrightarrow{\sim} \text{Hom}(\Pi', \bullet) ,$$

then by applying this functor to $\Pi' \in \mathcal{G}$, we obtain a homomorphism $\mu : \Pi \rightarrow \Pi'$ that corresponds to the identity automorphism $1_{\Pi'} \in \text{Hom}(\Pi', \Pi')$. Similarly, $1_{\Pi} \in \text{Hom}(\Pi, \Pi)$ corresponds to $\xi : \Pi' \rightarrow \Pi$. Because of the naturalness condition, for the homomorphism μ we have the commutative diagram

$$\begin{array}{ccccc} \Pi & \xrightarrow{1_{\Pi}} & \Pi & & \text{Hom}(\Pi, \Pi) & \xrightarrow{A_{\Pi}} & \text{Hom}(\Pi', \Pi) & & \Pi & \xleftarrow{\xi} & \Pi' \\ & & \mu \downarrow & & \downarrow & & \downarrow & & \mu \downarrow & & \\ & & \Pi' & & \text{Hom}(\Pi, \Pi') & \xrightarrow{A_{\Pi'}} & \text{Hom}(\Pi', \Pi') & & \Pi' & & \end{array}$$

that relates

$$\begin{array}{ccc} 1_{\Pi} & \longrightarrow & \xi \\ \downarrow & & \downarrow \\ \mu & \longrightarrow & 1_{\Pi'} \end{array} .$$

In particular, we have $\mu \circ \xi = 1_{\Pi'}$, hence $\Pi \cong \Pi'$. As a consequence of this faithfulness, we note that if two presentation functors are naturally isomorphic, then the two presented groups are actually isomorphic.

The above consideration suggests that we can use a particular presentation of a group Π to define the *volume* of the representation space $\text{Hom}(\Pi, G)$, which will be an invariant of the group Π , independent of its presentation. However, to carry out the calculation of the volume $|\text{Hom}(\Pi, G)|$, we need not only a presentation of Π , but also a more specific natural isomorphism between presentation functors. Notice that two presentations

$$(1.10) \quad \Pi_1 = \langle a_1, \dots, a_k \mid q_1 \dots, q_r \rangle \quad \text{and} \quad \Pi_2 = \langle b_1, \dots, b_k \mid s_1 \dots, s_r \rangle$$

define the same group if and only if for every a_i there is a word $a_i(b)$ of the generators $b_1, \dots, b_k$ and for every b_j there is a word $b_j(a)$ of $a_1, \dots, a_k$ such that for any group $G \in \mathcal{G}$, the maps a and b associated to these words are bijective and satisfy the commutativity

$$(1.11) \quad \begin{array}{ccccc} G^k & \xlongequal{\quad} & G^k & \xrightarrow{q} & G^r \\ a \uparrow \wr & & \wr \downarrow b & & \\ G^k & \xlongequal{\quad} & G^k & \xrightarrow[s]{} & G^r \end{array}$$

and

$$(1.12) \quad q^{-1}(1) = b^{-1}(s^{-1}(1)), \quad s^{-1}(1) = a^{-1}(q^{-1}(1)) .$$

In Section 2 we use this specific form of the natural isomorphism of the presentation functors to show the independence of the volume of the representation variety on the choice of presentation.

2. VOLUME OF REPRESENTATION VARIETIES

From now on we restrict our attention to the category of compact real analytic Lie groups. The representation space $\text{Hom}(\Pi, G)$ of

$$\Pi = \langle a_1, \dots, a_k \mid q_1, \dots, q_r \rangle$$

in a compact Lie group G is called the *representation variety*. Unless otherwise stated, we do not assume connectivity or semisimplicity for the compact Lie group G . Therefore, it can be in particular a finite group. The natural identification (1.8) makes $\text{Hom}(\Pi, G)$ a real analytic subvariety of G^r . In this section we define the volume of the representation variety using techniques from non-commutative harmonic analysis on compact Lie groups. Especially we use theory of distributions on a compact Lie group and character theory freely in Sections 2, 3 and 4. We refer to the summary presented in Section 5 for terminologies, definitions and properties of distributions necessary for our investigation.

We denote by dx a left and right invariant measure on G . We do not normalize the invariant measure, and call the quantity

$$|G| = \int_G dx$$

the volume of the group G . The invariant measure on the product group G^r is the product measure $dw_1 \cdots dw_r$. The δ -function on G^r is defined by

$$(2.1) \quad \delta_r(w_1, \dots, w_r) = \delta(w_1) \cdots \delta(w_r) .$$

For the presentation map

$$q : G^k \longrightarrow G^r$$

of (1.7), we associate the *volume distribution* $f_q(w)$ on G^r by

$$(2.2) \quad f_q(w) = |q^{-1}(w)| = \int_{G^k} \delta_r(q \cdot w^{-1}) dx_1 \cdots dx_k ,$$

where

$$q \cdot w^{-1} = (q_1(x)w_1^{-1}, \dots, q_r(x)w_r^{-1}) \in G^r$$

and $dx_1 \cdots dx_k$ is the product measure on G^k . The volume distribution is a Schwartz distribution on the group G^r characterized by the continuous linear functional associated with the push-forward measure

$$(2.3) \quad g(w) \longmapsto \int_{G^r} f_q(w) g(w) dw_1 \cdots dw_r = \int_{G^k} g(q(x_1, \dots, x_k)) dx_1 \cdots dx_k$$

for every $g(w) \in C^\infty(G^r)$. Sard's theorem tells us that the set of critical values of q has measure 0 in the target space with respect to the Lebesgue measure of G as a real analytic manifold, and hence also with respect to the Haar measure dx . If the distribution $f_q(w)$ is regular at $w = 1$ (see Section 5), then we define the *volume* of the representation variety by

$$(2.4) \quad |\mathrm{Hom}(\Pi, G)| = f_q(1) .$$

The total volume of $G^k = q^{-1}(G^r)$ with respect to the volume distribution is

$$\int_{G^r} f_q(w) dw_1 \cdots dw_r = \int_{G^k} \left(\int_{G^r} \delta_r(q \cdot w^{-1}) dw_1 \cdots dw_r \right) dx_1 \cdots dx_k = |G^k| .$$

Thus our definition (2.4) agrees with Fubini's volume if the presentation map q is a fiber bundle. Since evaluation of a distribution at a point does not usually make sense, we caution that the volume is defined *only when* $f_q(1)$ is regular. This imposes a strong condition on the presentation of Π .

Proposition 2.1. *If the presentation map $q : G^k \rightarrow G^r$ has $1 \in G^r$ as a regular value, then the volume distribution $f_q(w)$ is also regular at the identity element.*

Proof. The condition implies that the differential dq of the presentation map q has the maximal rank at every point of $q^{-1}(1)$, and that $q^{-1}(1)$ is a non-singular submanifold of G^k . Since G is compact, there is an open neighborhood U of $1 \in G^r$ such that

$$(2.5) \quad \phi : F \times U \xrightarrow{\sim} q^{-1}(U)$$

as a C^∞ manifold. Here $F = \mathrm{Hom}(\Pi, G) = q^{-1}(1)$ is a C^∞ manifold and the presentation map restricted to $q^{-1}(U)$ is the projection $F \times U \rightarrow U$. Let ω be the invariant volume form on G corresponding to the invariant measure dx . Denote by $p_j : G^k \rightarrow G$ the projection to the j -th component. Then

$$(2.6) \quad \omega^k \stackrel{\mathrm{def}}{=} p_1^* \omega \wedge \cdots \wedge p_k^* \omega$$

corresponds to the product measure $dx_1 \cdots dx_k$ on G^k . Similarly, we denote by ω^r the invariant volume form on the product group G^r corresponding to $dw_1 \cdots dw_r$. Notice that

the volume form $\phi^*\omega^k$ on $F \times U$ is a C^∞ form. Therefore, there is a C^∞ function $h(w)$ on U such that the integration of $\phi^*\omega^k$ along fiber associated with the fibration $F \times U \rightarrow U$ is given by

$$\int_{F \times \{w\}} \phi^*\omega^k = h(w) \omega^r .$$

Since the volume distribution $f_q(w)$ is the push-forward measure, we have the equality

$$(f_q|_U)(w) = h(w)$$

as a distribution. Therefore, the localization $f_q|_U$ of the volume distribution $f_q(w)$ to U is a smooth function. $\square$

Example 2.1. Consider a presentation

$$\Pi = \langle a, b \mid ab^{-1} = 1 \rangle .$$

The corresponding presentation map is

$$q : G \times G \ni (x, y) \longmapsto xy^{-1} \in G ,$$

hence $q^{-1}(1)$ is the diagonal of $G \times G$. The volume of the diagonal is calculated by

$$|\text{Hom}(\Pi, G)| = \int_{G^2} \delta(xy^{-1}) dx dy = |G| .$$

Notice that the same group is presented by $\langle a \rangle$. With this presentation, there is no relation, and the presentation map is the trivial map

$$G \longrightarrow \{1\} = G^0 .$$

The delta function on the trivial group is the constant $1 \in \mathbb{C}$, hence

$$|\text{Hom}(\langle a \rangle, G)| = \int_G dx = |G| .$$

For a compact Lie group G , we require that the maps a and b of the natural isomorphism of (1.11) and (1.12) are real analytic automorphisms of the real analytic manifold G^k .

Lemma 2.2. *Suppose*

$$b : G^k \ni x = (x_1, \dots, x_k) \longmapsto (b_1(x), \dots, b_k(x)) = b(x) \in G^k$$

is an analytic automorphism of the real analytic manifold G^k given by k words $b_1, \dots, b_k$ of $x_1, \dots, x_k \in G$ such that its inverse is also given by k words in a similar way. Then the volume form ω^k of G^k is invariant under the automorphism b up to a sign:

$$(2.7) \quad b^*\omega^k = \pm \omega^k .$$

Proof. Since the automorphism b maps $(1, \dots, 1) \in G^k$ to itself, the differential db at $(1, \dots, 1) \in G^k$ can be described in terms of the Lie algebra $\mathfrak{g}$ of G . Noticing that a word is a product of ± 1 powers of x_j 's, the differential

$$db : T_{(1, \dots, 1)} G^k = \mathfrak{g}^{\oplus k} \xrightarrow{\sim} \mathfrak{g}^{\oplus k} = T_{(1, \dots, 1)} G^k$$

is given by an invertible $k \times k$ matrix of integer entries $N \in GL(k, \mathbb{Z})$. Since $\det N = \pm 1$, the G -invariant volume form on $\mathfrak{g}^{\oplus k}$ is preserved by db up to the sign $(\det N)^{k \dim G}$.

Now notice that

$$(2.8) \quad \begin{aligned} b^*\omega^k &= b_1^*\omega \wedge \dots \wedge b_k^*\omega \\ \omega^k &= p_1^*\omega \wedge \dots \wedge p_k^*\omega , \end{aligned}$$

where $p_j : G^k \rightarrow G$ is the projection to the j -th component used in (2.6). To analyze the pull-back of ω by a word b_i , let us examine the pull-back of ω via the multiplication map

$$m : G^\nu \ni (y_1, \dots, y_\nu) \mapsto y_1 \cdots y_\nu \in G$$

of (1.5). Here ν is the length of b_j . Since ω is left and right multiplication invariant, the restriction of $m^*\omega$ to the ℓ -th factor of G^ν is ω itself. To be more precise, it is $p_\ell^*\omega$. If the ℓ -th letter of the word b_i is x_1 , then the restriction of

$$\iota_{b_i} : G^k \longrightarrow G^\nu$$

of (1.4) on the first factor of G^k into the ℓ -th factor of G^ν is the identity map. Thus $\iota_{b_i}^* m^* \omega$ restricted to the first factor of G^k is just an integer multiple of ω (or to be precise $p_1^*\omega$), and the coefficient is the number of times the letter x_1 appears in the word b_i plus $(-1)^{\dim G}$ for each appearance of x_1^{-1} in b_i . Define n_{ij} as the number of letter x_j in the word b_i minus the number of x_j^{-1} in b_i . Then the matrix $N = (n_{ij})$ is the same matrix N that represents the differential db . Therefore, $N \in GL(k, \mathbb{Z})$. Noticing the identity

$$b^* \omega^k = b_1^* \omega \wedge \cdots \wedge b_k^* \omega = \iota_{b_1}^* m^* \omega \wedge \cdots \wedge \iota_{b_k}^* m^* \omega ,$$

we obtain the desired formula

$$b^* \omega^k = N^* \omega^k = (\det N)^{k \dim G} \cdot \omega^k .$$

□

A consequence of this lemma is the independence of the volume $|\text{Hom}(\Pi, G)|$ on the choice of presentation among those with k generators and r relators.

Theorem 2.3. *Let G be a compact Lie group. Suppose we have two presentations (1.10) of the same group Π with the same k and r . If the presentation maps p and q have the regular value at $1 \in G^r$, then the volume of the representation variety of Π in G is independent of the presentation:*

$$(2.9) \quad f_q(1) = \int_{G^k} \delta_r(q(x_1, \dots, x_r)) dx_1 \cdots dx_k = \int_{G^k} \delta_r(s(y_1, \dots, y_k)) dy_1 \cdots dy_k = f_s(1) .$$

Proof. As in Proposition 2.1, there is an open neighborhood $U \ni 1$ of G^r and C^∞ maps ϕ and ψ such that

$$\begin{array}{ccccc} q^{-1}(U) & \xleftarrow[\phi]{\sim} & F \times U & \xrightarrow[\psi]{\sim} & (sb)^{-1}(U) \\ q \downarrow & & \downarrow \text{proj} & & \downarrow sb \\ U & \xlongequal{\quad} & U & \xlongequal{\quad} & U \end{array} ,$$

where $F = q^{-1}(1) = (sb)^{-1}(1) \subset G^k$. Note in particular that the restrictions of ϕ and ψ on $F \times \{1\}$ are the identity map. Let ω^k and ω^r denote the volume forms on G^k and G^r , respectively, defined by (2.6). Then the localizations of the volume distributions f_q and f_{sb} on U (see Section 5) are C^∞ functions and given by the integration along fiber:

$$\int_{F \times \{w\}} \phi^* \omega^k = (f_q|_U)(w) \omega^r , \quad \int_{F \times \{w\}} \psi^* \omega^k = (f_{sb}|_U)(w) \omega^r .$$

Choose a volume form ω_F on F and write

$$\phi^* \omega^k = h_q(z, w) \omega_F \wedge \omega^r \quad \text{and} \quad \psi^* \omega^k = h_{sb}(z, w) \omega_F \wedge \omega^r$$

using C^∞ functions h_q and h_{sb} on $F \times U$. Then the integration along fiber is given by

$$\begin{aligned} (f_q|_U)(w) \omega^r &= \int_{F \times \{w\}} \phi^* \omega^k = \left(\int_{F \times \{w\}} h_q(z, w) \omega_F \right) \omega^r \\ (f_{sb}|_U)(w) \omega^r &= \int_{F \times \{w\}} \psi^* \omega^k = \left(\int_{F \times \{w\}} h_{sb}(z, w) \omega_F \right) \omega^r . \end{aligned}$$

Since $h_q(z, 1) = h_{sb}(z, 1)$ as a function on $F \times \{1\}$, we obtain $f_q(1) = f_{sb}(1)$. Thus if we have $b^* \omega^k = \omega^k$, then $f_{sb}(1) = f_s(1)$, and the proof is completed. This invariance of the volume form ω^k under the automorphism b is assured by the previous lemma. The sign ± 1 in Lemma 2.2 is not our concern here because we always use the positive volume form for integration on G^k . $\square$

To establish the independence of the volume on the choice of presentation with different numbers of generators and relators, we first need its invariance under the Andrews-Curtis moves [1].

Definition 2.4. The *Andrews-Curtis moves* of a presentation (1.6) of a discrete group are the following six operations on the generators and relators:

- (1) Interchange q_1 and q_j for $j = 2, \dots, r$.
- (2) Replace q_1 with aq_1a^{-1} for an element $a \in \Pi$.
- (3) Replace q_1 with q_1^{-1} .
- (4) Replace q_1 with q_1q_2 .
- (5) Add another generator a and a relator $q_{r+1} = a$.
- (6) Delete a generator a and a relator a .

Theorem 2.5. *Let $\Pi = \langle a_1, \dots, a_k \mid q_1, \dots, q_r \rangle$ be a finite presentation of a discrete group and G a compact Lie group. If the volume distribution $f_q(w)$ is regular at $w = 1 \in G^r$ and hence the volume of the representation variety $f_q(1) = |\text{Hom}(\Pi, G)|$ is well-defined, then the volume is invariant under the Andrews-Curtis moves.*

Proof. The invariance under the first three moves follows immediately from (2.1) and (5.8). To show the invariance under (4), we note the following identity of δ -functions:

$$(2.10) \quad \delta_2(x, y) = \delta(x)\delta(y) = \delta(xy)\delta(y) .$$

This is because for every $g(x, y) \in C^\infty(G^2)$, we have

$$\begin{aligned} \int_{G^2} g(x, y) \delta(xy) \delta(y) dx dy &= \int_G g(y^{-1}, y) \delta(y) dy \\ &= g(1, 1) \\ &= \int_{G^2} g(x, y) \delta_2(x, y) dx dy . \end{aligned}$$

Therefore,

$$\delta(q_1)\delta(q_2) \cdots \delta(q_r) = \delta(q_1q_2)\delta(q_2) \cdots \delta(q_r) .$$

The invariance under (5) and (6) comes from the following commutative diagram

$$(2.11) \quad \begin{array}{ccc} G^k \times G & \xrightarrow{(q, id)} & G^r \times G \\ i_k \uparrow & & \uparrow i_r \\ G^k & \xrightarrow{q} & G^r \end{array} ,$$

where the vertical maps are the inclusions into the first G^k and G^r components, respectively, with $1 \in G$ in the last component. The invariance is a consequence of the identity

$$\int_{G^k \times G} \delta_r(q) \delta(x) dx_1 \cdots dx_k dx = \int_{G^k} \delta_r(q) dx_1 \cdots dx_k .$$

□

The Andrews-Curtis moves were introduced in [1] with the well-known conjecture stating that every presentation of the trivial group with the same number of generators and relations can be transformed into the standard presentation by a finite sequence of moves (1)–(6). The conjecture is still open to date. For three-manifold groups, the conjecture is expected to be true.

A three-manifold group $\Pi = \pi_1(M)$ of a closed oriented 3-manifold M always has a *balanced* presentation, meaning that the number of generators and relations are the same, given by a Heegaard splitting of M . If Π has a balanced presentation, then the expected dimension of the representation variety $\text{Hom}(\Pi, G)$ in a compact Lie group G is 0, and hence it consists of a finite set. The Casson invariant counts the number of elements of $\text{Hom}(\pi_1(M), SU(2))$ with sign. If indeed $\text{Hom}(\pi_1(M), G)$ is a finite set, then its cardinality $|\text{Hom}(\pi_1(M), G)|$ is a topological invariant of M . For example, Kuperberg [24] discussed this quantity for a finite group G . But often the representation variety contains higher dimensional exceptional components. If such a situation happens, then the volume distribution becomes singular at the identity element of G^r , and our theory does not provide any useful information.

We are now ready to prove that the volume of the representation variety is independent of the choice of presentation, if it is well-defined. More precisely, we have

Theorem 2.6. *Suppose a group Π has two presentations of the form*

$$(2.12) \quad \langle a_1, \dots, a_k \mid q_1, \dots, q_r \rangle$$

and

$$(2.13) \quad \langle b_1, \dots, b_\ell \mid s_1, \dots, s_t \rangle$$

such that

$$(2.14) \quad k - r = \ell - t ,$$

or in other words, the expected dimensions of $\text{Hom}(\Pi, G)$ through the two presentations agree. If the volume distributions f_q and f_s are regular at the identity element of the product groups G^r and G^t , respectively, then

$$|\text{Hom}(\Pi, G)| = f_q(1) = f_s(1) .$$

Proof. Without loss of generality, we can assume $k \leq \ell$. Using the Andrews-Curtis moves, we supply the new generators $a_{k+1}, \dots, a_\ell$ and relators

$$q_{r+1} = a_{k+1}, \dots, q_{r+\ell-k} = a_\ell$$

to the first presentation (2.12) so that we have

$$(2.15) \quad \Pi = \langle a_1, \dots, a_\ell \mid q_1, \dots, q_{r+\ell-k} \rangle .$$

From (2.14), we know $r + \ell - k = t$. Therefore, the two presentations (2.15) and (2.13) have the same numbers of generators and relators. If $f_q(w)$ is regular at $w = 1 \in G^r$, then the volume distribution of the enlarged presentation (2.15) is also regular at $1 \in G^\ell$. Therefore, from Theorem 2.3, we conclude that

$$f_q(1) = f_s(1) .$$

□

For a one-relation group such as a surface group, character theory of compact Lie groups provides a computational tool. In the next two sections, we calculate the volume distribution for every surface group in terms of irreducible characters of G .

Lemma 2.7. *Let $\Pi = \langle a_1, \dots, a_k \mid q \rangle$ be a one-relation group. Then for every compact Lie group G , the volume distribution (2.2) is a class distribution on G .*

Proof. For $y \in G$ and $g(w) \in C^\infty(G)$, we have

$$\begin{aligned} \int_G f_q(ywy^{-1})g(w)dw &= \int_G f_q(w)g(y^{-1}wy)dw \\ &= \int_{G^k} g(y^{-1}q(x_1, \dots, x_k)y)dx_1 \cdots dx_k \\ &= \int_{G^k} g(q(y^{-1}x_1y, \dots, y^{-1}x_ky))dx_1 \cdots dx_k \\ &= \int_{G^k} g(q(x_1, \dots, x_k))dx_1 \cdots dx_k \\ &= \int_G f_q(w)g(w)dw \end{aligned}$$

because of the left and right invariance of the Haar measure. □

Since the irreducible characters are real analytic functions on G and form an orthonormal basis for the L^2 class functions on G , we have

Lemma 2.8. *The character expansion of the class distribution $f_q(w)$ on G is given by*

$$(2.16) \quad f_q(w) = \frac{1}{|G|} \sum_{\lambda \in \hat{G}} \left(\int_{G^k} \bar{\chi}_\lambda(q(x_1, \dots, x_k))dx_1 \cdots dx_k \right) \chi_\lambda(w) .$$

Notice that the integral in (2.16) is convergent and well-defined, and the infinite sum is convergent with respect to the strong topology of $\mathcal{D}'(G)$. Thus the regularity question of $f_q(w)$ can be answered by studying the convergence problem of the infinite sum over all irreducible representations. For example, if the sum of (2.16) is convergent with respect to the sup norm on G , then f_q is a C^∞ function on G (see Section 5).

3. REPRESENTATION VARIETIES OF ORIENTABLE SURFACE GROUPS

In this section we calculate the volume distribution of the surface group $\pi_1(S)$ in G and the volume $|\text{Hom}(\pi_1(S), G)|$ of the representation variety for a closed orientable surface S and a compact Lie group G . In this and the next section, the statements and the argument of proof are actually almost identical to that of the classical papers by Frobenius and Schur [11, 12]. Because of the way we define the volume in the previous section, the same argument simultaneously works for both finite groups and compact Lie groups.

Let S be a closed orientable surface of genus $g \geq 1$. Its fundamental group is given by

$$(3.1) \quad \pi_1(S) = \langle a_1, b_1, \dots, a_g, b_g \mid [a_1, b_1] \cdots [a_g, b_g] \rangle ,$$

where $[a, b] = aba^{-1}b^{-1}$ is the group commutator of a and b . The presentation map is a real analytic map

$$(3.2) \quad q_g : G^{2g} \ni (x_1, y_1, \dots, x_g, y_g) \longmapsto [x_1, y_1] \cdots [x_g, y_g] \in G .$$

The representation variety of $\pi_1(S)$ in G is a real analytic subvariety $\text{Hom}(\pi_1(S), G) = q_g^{-1}(1)$ of G^{2g} . For every $g \geq 1$, we denote by f_g the volume distribution

$$(3.3) \quad f_g(w) = f_{q_g}(w) = \int_{G^{2g}} \delta([x_1, y_1] \cdots [x_g, y_g] w^{-1}) dx_1 dy_1 \cdots dx_g dy_g .$$

By Lemma 2.7, f_g is a class distribution. Our purpose is to calculate its character expansion. From the convolution property of the delta function

$$\delta(q_{g+h} w^{-1}) = \int_G \delta(q_g w^{-1} u^{-1}) \delta(u q_h) du ,$$

we have

$$(3.4) \quad f_{g+h} = f_g * f_h = f_h * f_g$$

for every $g, h \geq 1$. More precisely, consider the breakdown of q_{g+h} :

$$\begin{array}{ccc} G^{2g} \times G^{2h} & \ni & (x_1, y_1, \dots, x_{g+h}, y_{g+h}) \\ \downarrow (q_g \times q_h) & & \downarrow \\ G \times G & \ni & ([x_1, y_1] \cdots [x_g, y_g], [x_{g+1}, y_{g+1}] \cdots [x_{g+h}, y_{g+h}]) \\ \downarrow \text{multiplication} & & \downarrow \\ G & \ni & [x_1, y_1] \cdots [x_{g+h}, y_{g+h}] . \end{array}$$

The distribution f_{g+h} is the push-forward of the product measure on G^{2g+2h} via q_{g+h} , which factors into the convolution product according to (3.4). Here we have used the fact that the convolution of two distributions f_g and f_h corresponds to the push-forward measure via the multiplication map $G \times G \rightarrow G$. In particular, f_g is the g -th convolution power of f_1 :

$$f_g = \overbrace{f_1 * \cdots * f_1}^{g\text{-times}} .$$

Therefore, it suffices to find the character expansion of f_1 . The following formula and its proof is essentially due to Frobenius [11] published in 1896. We translate his argument from a finite group to a compact Lie group using the Dirac delta function and integration over the group.

Proposition 3.1. *As a distribution on G , we have the following strongly convergent character expansion formula in $\mathcal{D}'(G)$:*

$$(3.5) \quad f_1(w) = \int_{G^2} \delta(xy x^{-1} y^{-1} w^{-1}) dx dy = \sum_{\lambda \in \hat{G}} \frac{|G|}{\dim \lambda} \chi_\lambda(w) .$$

Proof. We begin with the convolution formula for the push-forward measure

$$\delta(xy x^{-1} y^{-1} w^{-1}) = \int_G \delta(x(wu^{-1})^{-1}) \delta(yx^{-1} y^{-1} u^{-1}) du .$$

We use the class distribution of (5.16) and its character expansion (5.17) here to obtain

$$\eta_{x^{-1}}(u) = \int_G \delta(yx^{-1} y^{-1} u^{-1}) dy = \sum_{\lambda \in \hat{G}} \bar{\chi}_\lambda(x^{-1}) \chi_\lambda(u) = \sum_{\lambda \in \hat{G}} \chi_\lambda(x) \chi_\lambda(u) .$$

Then we have

$$\begin{aligned} f_1(w) &= \int_{G^2} \delta(xy x^{-1} y^{-1} w^{-1}) dx dy \\ &= \int_{G^3} \delta(x(wu^{-1})^{-1}) \delta(yx^{-1} y^{-1} u^{-1}) du dx dy \\ &= \int_{G^2} \delta(x(wu^{-1})^{-1}) \eta_{x^{-1}}(u) du dx \\ &= \sum_{\lambda \in \hat{G}} \int_{G^2} \delta(x(wu^{-1})^{-1}) \chi_\lambda(x) \chi_\lambda(u) du dx \\ &= \sum_{\lambda \in \hat{G}} \int_G \chi_\lambda(wu^{-1}) \chi_\lambda(u) du \\ &= \sum_{\lambda \in \hat{G}} \frac{|G|}{\dim \lambda} \chi_\lambda(w) . \end{aligned}$$

□

Taking the g -th convolution power of f_1 , we obtain

Theorem 3.2. *For every $g \geq 1$, the character expansion of the volume distribution $f_g = f_{q_g}$ is given by*

$$(3.6) \quad f_g(w) = f_{q_g}(w) = \sum_{\lambda \in \hat{G}} \left(\frac{|G|}{\dim \lambda} \right)^{2g-1} \chi_\lambda(w) .$$

If the sum of the RHS converges with respect to the sup norm on G , then the value of $f_g(w)$ gives the volume $|q_g^{-1}(w)|$, and in particular, we have

$$(3.7) \quad |G|^{\chi(S)-1} |\mathrm{Hom}(\pi_1(S), G)| = \sum_{\lambda \in \hat{G}} (\dim \lambda)^{\chi(S)} .$$

Conversely, if the RHS of (3.7) is convergent, then the character expansion (3.6) is uniformly and absolutely convergent and the volume distribution is a C^∞ class function.

Remark. Except for the finite group case, the RHS of (3.7) never converges for $g = 0$ and $g = 1$. The Weyl dimension formula gives a lower bound on the genus g for the series to converge. For example, if $G = SU(n)$, then the series (3.7) converges for every $g \geq 2$.

We note that the argument works without any modification for a finite group. If G is a finite group, then there is a canonical choice of the Haar measure: a discrete measure of uniform weight 1 for each element. With this choice, the volume $|G|$ is the order of the group and the delta function is the characteristic function that takes value 1 at the identity element and 0 everywhere else. There is no difference between distributions and differentiable functions in this case, and there is no problem of convergence in the character expansion. Thus $f_g(w)$ of (3.6) is a function with a well-defined value for every $w \in G$. In particular, the expression of $f_g(1)$ for $g \geq 1$ recovers Mednykh's formula [29] (Theorem 0.3). For a surface of genus 0, the formula reduces to the classical formula (0.8) and remains true.

For the regularity of the distribution of $f_g(w)$ at $w = 1$, we have the following:

Proposition 3.3. *Let S be a closed orientable surface of genus g and G a compact connected semisimple Lie group of dimension n . If $g \geq n$, then the volume distribution of (3.3) is regular at $w = 1$.*

Proof. Since G is semisimple, its Lie algebra $\mathfrak{g}$ satisfies

$$(3.8) \quad D(\mathfrak{g}) = [\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}.$$

This means every element of G near the identity is expressible as a finite product of commutators. Let us analyze the map

$$q_1 : G \times G \ni (x, y) \longmapsto xyx^{-1}y^{-1} \in G$$

near the identity element. Put $x = e^{tA}$ and $y = e^{sB}$ for some $A, B \in \mathfrak{g}$. Then from the Campbell-Hausdorff formula we have

$$\begin{aligned} xyx^{-1}y^{-1} &= e^{tA+sB+\frac{ts}{2}[A,B]+\dots} e^{-tA-sB+\frac{ts}{2}[A,B]+\dots} \\ &= 1 + ts[A, B] + \text{higher} . \end{aligned}$$

Therefore, for a fixed $y = e^B$, the tangent line at the identity of the image of the one-parameter subgroup e^{tA} via q_1 is given by $t[A, B] \in \mathfrak{g}$.

Now let $(A_1, B_1, \dots, A_g, B_g) \in \mathfrak{g}^{2g}$ be an arbitrary $2g$ -tuple of Lie algebra elements. The differential of the map $q_g : G^{2g} \rightarrow G$ at $(1, \dots, 1)$ of G^{2g} is

$$(3.9) \quad dq_g : \mathfrak{g}^{2g} \ni (A_1, B_1, \dots, A_g, B_g) \longmapsto [A_1, B_1] + \dots + [A_g, B_g] \in \mathfrak{g}.$$

If $g \geq n = \dim G$, then dq_g is surjective because of (3.8). This implies that the map q_g is regular at $(1, \dots, 1)$.

Let $(a_1, b_1, \dots, a_g, b_g) \in q_g^{-1}(1)$ be an arbitrary point of the inverse image of 1 via q_g . Since the exponential map is surjective, we can write $a_j = e^{X_j}$ and $b_j = e^{Y_j}$ for $j = 1, \dots, g$. Then again by the Campbell-Hausdorff formula, the differential dq_g at $(a_1, b_1, \dots, a_g, b_g)$ gives for $(A_1, B_1, \dots, A_g, B_g) \in \mathfrak{g}^{2g}$ the value

$$\sum_{j=1}^g ([X_j, B_j] + [A_j, Y_j] + [A_j, B_j]) + \dots ,$$

where we used the fact that $(a_1, b_1, \dots, a_g, b_g)$ is in the inverse image of 1. This is a surjective map onto $\mathfrak{g}$. Therefore, q_g is locally a fibration in a neighborhood of the identity, and hence the volume distribution $f_g(w)$ is regular at $w = 1$ by Proposition 2.1. $\square$

- Remark.* (1) The semisimplicity condition of G is necessary. Indeed, if G has a center Z of positive dimension, then $Z \times G$ is in $q_1^{-1}(1)$, which has a larger dimension than the expected dimension of the inverse image. Thus the map q_g has critical value at 1, and hence $f_g(w)$ may not be regular at $w = 1$. In particular, the volume distribution is always singular at the identity for any compact abelian group of positive dimension.
- (2) Even though $f_g(1)$ is regular, it does not imply that the series of the RHS of (3.7) is convergent.

Example 3.1. Let us consider the case $G = SU(2)$. Since $\hat{G} = \mathbb{N}$, we identify λ with a positive integer n , which is the dimension of λ . The character expansion (3.6) is uniformly and absolutely convergent for $g \geq 2$, and we have for every closed orientable surface S of $g \geq 2$,

$$\frac{|\text{Hom}(\pi_1(S), SU(2))|}{|SU(2)|^{2g-1}} = \zeta(2g-2) .$$

4. REPRESENTATION VARIETIES OF NON-ORIENTABLE SURFACE GROUPS

Let us now turn our attention to the case of a non-orientable surface S . We recall that every closed non-orientable surface is obtained by removing $k \geq 1$ disjoint disks from a sphere S^2 and gluing a cross-cap to each hole. The number k is the *cross-cap genus* of S , and its Euler characteristic is given by $\chi(S) = 2 - k$. Since the fundamental group of a non-orientable surface of cross-cap genus k is given by

$$(4.1) \quad \pi_1(S) = \langle a_1, \dots, a_k \mid a_1^2 \cdots a_k^2 \rangle ,$$

we use the presentation map

$$(4.2) \quad r_k : G^k \ni (x_1, \dots, x_k) \longmapsto x_1^2 \cdots x_k^2 \in G .$$

Our aim is to compute the volume distribution

$$(4.3) \quad \begin{aligned} h_k(w) = f_{r_k}(w) &= |r_k^{-1}(w)| = \int_{G^k} \delta(r_k(x_1, \dots, x_k) \cdot w^{-1}) dx_1 \cdots dx_k \\ &= \int_{G^k} \delta(x_1^2 \cdots x_k^2 w^{-1}) dx_1 \cdots dx_k . \end{aligned}$$

Using the same technique of Section 3, it follows from the convolution property of the δ -function that

$$h_k = \overbrace{h_1 * \cdots * h_1}^{k\text{-times}} .$$

Proposition 4.1. *The character expansion of the class distribution h_1 is given by*

$$(4.4) \quad h_1(w) = \int_G \delta(x^2 w^{-1}) dx = \sum_{\lambda \in \hat{G}} \phi(\lambda) \chi_\lambda(w) ,$$

where

$$(4.5) \quad \phi(\lambda) = \frac{1}{|G|} \int_G \chi_\lambda(x^2) dx$$

is the Frobenius-Schur indicator of [12].

Proof. Let $h_1(w) = \sum_{\lambda} a_{\lambda} \chi_{\lambda}(w)$ be the expansion in terms of irreducible characters with respect to the strong convergence in $\mathcal{D}'(G)$. By definition, we have

$$\begin{aligned} a_{\lambda} &= \frac{1}{|G|} \int_G h_1(w) \overline{\chi}_{\lambda}(w) dw \\ &= \frac{1}{|G|} \int_G \int_G \delta(x^2 w^{-1}) \overline{\chi}_{\lambda}(w) dw dx \\ &= \frac{1}{|G|} \int_G \overline{\chi}_{\lambda}(x^2) dx \\ &= \frac{1}{|G|} \int_G \chi_{\lambda}(x^2) dx . \end{aligned}$$

□

Remark. Since the presentation map r_1 has the same domain and the range, if the space $r_1^{-1}(1)$ of involutions of G has positive dimension, then the map $r_1 : G \ni x \mapsto x^2 \in G$ does not have regular value at $1 \in G$. Let G be a compact connected Lie group of rank r and T a maximal torus of G . A compact torus of dimension r has exactly 2^r involutions. The conjugacy class of each of these involutions has positive dimension unless the involution is central. Thus the groups $SO(n)$ and $SU(n)$ for $n \geq 3$ and $Sp(n)$ for $n \geq 2$ all have large space of involutions with positive dimensions. Therefore, the volume $h_1(1)$ is ill-defined for these groups.

Taking the k -th convolution power of h_1 , we have

Theorem 4.2. *Let S be a closed non-orientable surface of cross-cap genus $k \geq 1$ and G an arbitrary compact Lie group. Then the character expansion of the volume distribution is given by*

$$(4.6) \quad h_k(w) = |r_k^{-1}(w)| = \sum_{\lambda \in \hat{G}_1} \left(\frac{|G|}{\dim \lambda} \right)^{k-1} \chi_{\lambda}(w) - \sum_{\lambda \in \hat{G}_4} \left(-\frac{|G|}{\dim \lambda} \right)^{k-1} \chi_{\lambda}(w) .$$

If the character sum of the RHS is absolutely convergent with respect to the sup norm on G , then the value gives the volume $|r_k^{-1}(w)|$. In particular, we have

$$(4.7) \quad |G|^{\chi(S)-1} |\text{Hom}(\pi_1(S), G)| = \sum_{\lambda \in \hat{G}_1} (\dim \lambda)^{\chi(S)} + \sum_{\lambda \in \hat{G}_4} (-\dim \lambda)^{\chi(S)} .$$

Conversely, if the RHS of (4.7) is absolutely convergent, then (4.6) is uniformly and absolutely convergent and the volume distribution h_k is a C^{∞} class function.

Proof. All we need is to notice the value of the Frobenius-Schur indicator (0.6). □

Example 4.1. Consider an n -dimensional torus $G = T^n$. Since the only real irreducible representation is the trivial representation, $\hat{G}_1 = \{1\}$. Let $S^1 = \{e^{i\theta} \mid 0 \leq \theta < 2\pi\}$ and choose an invariant measure $d\theta$ on it. The δ -function on the torus is the product of

δ -functions on S^1 . Thus we have

$$\begin{aligned} h_1(w) &= \int_{T^n} \delta(2\theta_1 - w_1) \cdots \delta(2\theta_n - w_n) d\theta_1 \cdots d\theta_n \\ &= \left(\int_0^{2\pi} \delta(2\theta) d\theta \right)^n \\ &= \left(\frac{1}{2} \int_0^{4\pi} \delta(\theta) d\theta \right)^n \\ &= 1. \end{aligned}$$

Therefore, $h_k(w) = |T^n|^{k-1}$, and hence $|\text{Hom}(\pi_1(S), T^n)| = |T^n|^{1-\chi(S)}$. The inverse image $r_k^{-1}(w)$ consists of $(x_1, \dots, x_k) \in (T^n)^k$ such that

$$(4.8) \quad x_k^2 = x_{k-1}^{-2} \cdots x_1^{-2} w.$$

Thus we can choose arbitrary $(x_1, \dots, x_{k-1}) \in (T^n)^{k-1}$, and for this choice, there are still 2^n solutions of (4.8). Each piece has the volume equal to $|T^n|^{k-1}/2^n$. Therefore, the total volume is $|T^n|^{k-1}$, in agreement with our computation.

Example 4.2. For $G = SU(2)$, we have a more interesting relation. We note that $\hat{G}_1$ consists of odd integers and $\hat{G}_4$ even integers. Therefore, the series (4.7) is absolutely convergent for $k \geq 4$. Thus for a closed non-orientable surface of cross-cap genus $k \geq 4$, we have

$$\begin{aligned} \frac{|\text{Hom}(\pi_1(S), SU(2))|}{|SU(2)|^{k-1}} &= \sum_{n=1}^{\infty} (2n-1)^{2-k} + (-1)^{2-k} \sum_{n=1}^{\infty} (2n)^{2-k} \\ &= \begin{cases} \zeta(k-2) & k \text{ is even} \\ \left(1 - \frac{1}{2^{k-3}}\right) \zeta(k-2) & k \text{ is odd} \end{cases}. \end{aligned}$$

Although the series is not absolutely convergent, we expect that the value for $k = 3$ is $\log(2)$.

In the same way as in Proposition 3.3, we have a general regularity condition of the volume distribution $h_k(w)$ for a compact semisimple Lie group.

Proposition 4.3. *Let G be a compact connected semisimple Lie group of dimension n and S a closed non-orientable surface of cross-cap genus $k \geq 1$. Define*

$$m = \left\lceil \frac{k-1}{2} \right\rceil = \begin{cases} (k-1)/2 & k \text{ is odd} \\ (k-2)/2 & k \text{ is even.} \end{cases}$$

Then the volume distribution $h_k(w)$ is regular at $w = 1$ if $m \geq n$.

Proof. The argument is based on another presentation of the fundamental group containing m commutator products

$$\pi_1(S) = \langle a_1, b_1, \dots, a_m, b_m, c \mid [a_1, b_1] \cdots [a_m, b_m] c^2 \rangle$$

for odd k and

$$\pi_1(S) = \langle a_1, b_1, \dots, a_m, b_m, c_1, c_2 \mid [a_1, b_1] \cdots [a_m, b_m] c_1^2 c_2^2 \rangle$$

for even k . First we notice that the substitution

$$(4.9) \quad \begin{cases} \alpha = abc \\ \beta = c^{-1}b^{-1}a^{-1}c^{-1}a^{-1}c \\ \gamma = c^{-1}ac^2, \end{cases}$$

yields the equality

$$(4.10) \quad [a, b] c^2 = \alpha^2 \beta^2 \gamma^2.$$

The inverse transformation of (4.9) is given by

$$(4.11) \quad \begin{cases} a = \alpha\beta\gamma\beta^{-1}\alpha^{-1}\gamma^{-1}\beta^{-1}\alpha^{-1} \\ b = \alpha\beta\gamma\alpha\beta\gamma^{-1}\beta^{-1}\gamma^{-1}\beta^{-1}\alpha^{-1} \\ c = \alpha\beta\gamma. \end{cases}$$

Therefore, the map

$$h : G^3 \ni (x, y, z) \longmapsto (xyz, z^{-1}y^{-1}x^{-1}z^{-1}x^{-1}z, z^{-1}xz^2) = (u, v, w) \in G^3$$

defined by (4.9) is an analytic automorphism of the manifold G^3 . Since both h and h^{-1} are given by words, from Lemma 2.2 we conclude

$$(4.12) \quad h^* \omega^3 = \omega^3.$$

Indeed, the integer matrix N representing the differential dh (see proof of Lemma 2.2) is

$$N = \begin{bmatrix} 1 & 1 & 1 \\ -2 & -1 & -1 \\ 1 & 0 & 1 \end{bmatrix}$$

with $\det N = 1$. Let

$$s_3 : G^3 \ni (x, y, z) \longmapsto [x, y] z^2 \in G$$

be the presentation map with respect to the relator $[a, b] c^2$, and

$$r_3 : G^3 \ni (u, v, w) \longmapsto u^2 v^2 w^2 \in G$$

the presentation map of $\alpha^2 \beta^2 \gamma^2$. Then (4.10) implies that the following diagram commutes:

$$\begin{array}{ccc} G^3 & \xrightarrow{s_3} & G \\ h \downarrow \wr & & \parallel \\ G^3 & \xrightarrow{r_3} & G \end{array}.$$

Together with the invariance of the volume form (4.12), we obtain

$$f_{s_3}(w) = f_{r_3}(w) = h_3(w)$$

as a distribution on G .

We can apply the above procedure based on the transformation (4.9) repeatedly to change the relator

$$[a_1, b_1] \cdots [a_m, b_m] c^2 \quad \text{to} \quad \alpha_1^2 \beta_1^2 \cdots \alpha_m^2 \beta_m^2 \gamma^2,$$

and

$$[a_1, b_1] \cdots [a_m, b_m] c_1^2 c_2^2 \quad \text{to} \quad \alpha_1^2 \beta_1^2 \cdots \alpha_m^2 \beta_m^2 \gamma_1^2 \gamma_2^2.$$

At each step the volume form on G^k is kept invariant, and we have the equality of the volume distributions

$$(4.13) \quad f_{s_k}(w) = h_k(w) ,$$

where $s_k : G^k \rightarrow G$ is the presentation map associated with the relator containing m commutators. The regularity of $f_{s_k}(w)$ at $w = 1 \in G$ can be established in the same way as in Proposition 3.3. The regularity of $h_k(w)$ follows from (4.13). $\square$

5. HARMONIC ANALYSIS AND IRREDUCIBLE REPRESENTATIONS OF A COMPACT LIE GROUP

In this section we briefly review the necessary accounts from the character theory of compact Lie groups and non-commutative harmonic analysis that are used in the main part of this paper. The important results we need are Schur's orthogonality relations of irreducible characters in terms of the convolution product, definition of the δ -function on a compact Lie group, and expansion of a class distribution in terms of irreducible characters. We refer to [6, 13, 49, 50, 51] for more detail.

Let G be a compact Lie group. We choose a left and right invariant, but not necessarily normalized, measure dx (Haar measure) on G that satisfies

$$(5.1) \quad \int_G g(x)dx = \int_G g(yx)dx = \int_G g(xy)dx = \int_G g(x^{-1})dx ,$$

where $g(x)$ is a smooth function on G and $y \in G$. The space of *distributions* $\mathcal{D}'(G)$ is the topological linear space consisting of all continuous linear functions on $C^\infty(G)$ with respect to the Fréchet topology of $C^\infty(G)$ [13, 51]. The left regular representation of G on $\mathcal{D}'(G)$ is defined by

$$(5.2) \quad \rho_L(w) : \mathcal{D}'(G) \ni f(x) \mapsto f(w^{-1}x) \in \mathcal{D}'(G) ,$$

where the distribution $f(w^{-1}x)$ is characterized by

$$(5.3) \quad \int_G f(w^{-1}x)g(x)dx = \int_G f(x)g(wx)dx$$

for every $g(x) \in C^\infty(G)$ using the invariance (5.1) of dx . Similarly, the right regular representation is defined by

$$(5.4) \quad \rho_R(w) : \mathcal{D}'(G) \ni f(x) \mapsto f(xw) \in \mathcal{D}'(G) .$$

Reflecting the group structure of the compact space G , $\mathcal{D}'(G)$ has the structure of an associative algebra over $\mathbb{C}$ whose multiplication is the *convolution product* of distributions f_1 and f_2 defined by

$$(5.5) \quad (f_1 * f_2)(x) = \int_G f_1(xw^{-1})f_2(w)dw .$$

A distribution $f(x) \in \mathcal{D}'(G)$ is a *class distribution* if

$$(5.6) \quad \rho_L(w^{-1}) \cdot f(x) = \rho_R(w) \cdot f(x) ,$$

or more conveniently, when $f(w^{-1}xw) = f(x)$ as a distribution for every $w \in G$. Class distributions form the center of the convolution algebra $\mathcal{D}'(G)$ because of the invariance of the Haar measure. Indeed, for every smooth function $g(x) \in C^\infty(G)$, the condition

$$(f * g)(x) = \int_G f(xw^{-1})g(w)dw = \int_G g(xu^{-1})f(u)du = (g * f)(x)$$

for $f(x) \in \mathcal{D}'(G)$ is equivalent to (5.6) because

$$\int_G g(xu^{-1})f(u)du = \int_G f(w^{-1}x)g(w)dw .$$

The Dirac δ -function on G is defined by

$$(5.7) \quad \int_G \delta(x)g(x)dx = g(1) , \quad g(x) \in C^\infty(G) .$$

This definition, together with the left and right invariance of the Haar measure dx , implies the following formulas for an arbitrary $y \in G$:

$$(5.8) \quad \begin{aligned} \int_G \delta(xy^{-1})f(x)dx &= f(y) , \quad f(x) \in \mathcal{D}'(G) \\ \delta(yxy^{-1}) &= \delta(x) = \delta(x^{-1}) \\ \delta(xy) &= \delta(yx) . \end{aligned}$$

Therefore, the δ -function is a class distribution and serves as the identity element of the convolution algebra $(\mathcal{D}'(G), *)$. In particular, we have $\delta * \delta = \delta$.

The Hermitian inner product of L^2 functions f_1 and f_2 on G is given by

$$(5.9) \quad \langle f_1, f_2 \rangle = \frac{1}{|G|} \int_G f_1(x) \overline{f_2(x)} dx .$$

Peter-Weyl theory provides a Hilbert basis for $L^2(G)$. The L^2 class functions on G form a Hilbert space with a Hilbert basis

$$(5.10) \quad \{\chi_\lambda \mid \lambda \in \hat{G}\} ,$$

where $\hat{G}$ is the set of equivalence classes of complex irreducible representations of G and χ_λ the character of $\lambda \in \hat{G}$. The irreducible characters satisfy *Schur's orthogonality relations*

$$(5.11) \quad \langle \chi_\lambda, \chi_\mu \rangle = \delta_{\lambda\mu}$$

and

$$(5.12) \quad \chi_\lambda * \chi_\mu = \frac{|G|}{\dim \lambda} \delta_{\lambda\mu} \chi_\lambda ,$$

where $\dim \lambda$ is the dimension of the representation $\lambda \in \hat{G}$.

Since G is compact, every finite dimensional representation has a natural G -invariant Hermitian inner product so that the representation is unitary. In particular, we have

$$(5.13) \quad |\chi_\lambda(x)| \leq |\chi_\lambda(1)| = \dim \lambda \quad \text{for every } x \in G .$$

Therefore, if a sequence $a_\lambda \in \mathbb{C}$ satisfies the convergence condition

$$(5.14) \quad \sup_{x \in G} \left| \sum_{\lambda \in \hat{G}} a_\lambda \chi_\lambda(x) \right| \leq \sum_{\lambda \in \hat{G}} |a_\lambda| \dim \lambda < \infty ,$$

then the series $\sum_{\lambda} a_{\lambda} \chi_{\lambda}$ is uniformly and absolutely convergent and defines a C^{∞} class function on G . Note that every L^2 class function on G can be expanded into a series of the form $\sum_{\lambda} a_{\lambda} \chi_{\lambda}$ with the convergence condition $\sum_{\lambda} |a_{\lambda}|^2 < \infty$. (If G is a compact connected abelian group, i.e., a compact torus, then the expansion in terms of irreducible characters is the Fourier series expansion.) In the same way, a class distributions on G is expanded into a series in χ_{λ} that is not necessarily absolutely convergent. For example, the δ -function has an expansion

$$(5.15) \quad \delta(x) = \frac{1}{|G|} \sum_{\lambda \in \hat{G}} \dim \lambda \cdot \chi_{\lambda}(x) .$$

This infinite sum is convergent with respect to the strong topology of the topological linear space $\mathcal{D}'(G)$, and gives an alternative definition of the delta function. We note that the convolution relation $\delta * \delta = \delta$ is equivalent to Schur's orthogonality relation (5.12).

The following family of class distributions parametrised by $x \in G$ are used in Section 3:

$$(5.16) \quad \eta_x(w) = \int_G \delta(yxy^{-1}w^{-1})dy .$$

It is the distribution corresponding to the linear functional

$$g(w) \mapsto \int_G g(yxy^{-1})dy$$

for every smooth function $g(w)$. It is a class distribution because for every $u \in G$,

$$\int_G g(yxy^{-1})dy = \int_G g(uyxy^{-1}u^{-1})d(uy) = \int_G g(u(yxy^{-1})u^{-1})dy .$$

The character expansion of $\eta_x(w)$ as a distribution in w has the following simple form:

$$(5.17) \quad \eta_x(w) = \sum_{\lambda \in \hat{G}} \bar{\chi}_{\lambda}(x) \chi_{\lambda}(w) .$$

Distributions cannot be evaluated in a meaningful way in general. For example, $\delta(1)$ is not defined. Similarly, restriction of a distribution defined on G to a closed subset is generally meaningless. However, a distribution can always be localized to any open subset of G . Let $U \subset G$ be an open subset and $\mathcal{D}(U)$ the Fréchet space of C^{∞} functions on U with compact support. The *localization* of a distribution on G to U ,

$$(5.18) \quad \mathcal{D}'(G) \ni f \mapsto f|_U \in \mathcal{D}'(U) ,$$

is defined by

$$\int_U (f|_U)(x)g(x)dx = \int_G f(x)\tilde{g}(x)dx$$

for every $g(x) \in \mathcal{D}(U)$, where $\tilde{g}(x)$ is the extension of $g(x)$ as a C^{∞} function on G satisfying

$$\tilde{g}(x) = \begin{cases} g(x) & x \in U \\ 0 & x \notin U \end{cases} .$$

Since $C^{\infty}(U) \subset \mathcal{D}'(U)$, we can compare $f|_U$ with a smooth function on U . A distribution $f \in \mathcal{D}'(G)$ is said to be *regular* at $w \in G$ if there is an open neighborhood $U \ni w$ such that the localization $f|_U$ of f is a C^{∞} function on U . If f is regular at $w \in G$, then the value $f(w) \in \mathbb{C}$ of f at w is well-defined.

6. REPRESENTATION VARIETIES AND CELL-DECOMPOSITIONS OF A SURFACE

We have observed that for every closed surface S and a compact Lie groups G , the quantity $|G|^{\chi(S)-1}|\text{Hom}(\pi_1(S), G)|$ is a natural object that does not depend on the choice of an invariant measure on G . In a recent work [35, 36], it has been noted that the same quantity for a finite group appears in the graphical expansion of a certain integral over the group algebra of the finite group. In this section we present a *heuristic* argument that relates the volume of the representation variety of a surface group to each term of the Feynman diagram expansion of a hypothetical “integral” over the group algebra of compact Lie groups. The argument is based on an alternative definition of the volume of the representation variety of a surface group in terms of a cell-decomposition of the surface.

The 1-skelton of a cell-decomposition of a closed connected surface is a *graph* drawn on the surface. Such graphs are known as *maps*, *fatgraphs*, *ribbon graphs*, *dessins d’enfants*, or *Möbius graphs*, depending on the context of study. Fatgraphs and ribbon graphs are used effectively in the study of topological properties of moduli spaces of pointed Riemann surfaces and related topics (see for example, [4, 18, 19, 22, 39, 40, 41, 42, 53]). Dessins d’enfants appear in the context of algebraic curves defined over the field of algebraic numbers (cf. [3, 17, 32, 33, 45, 46]). Maps are drawn on a topological surface and are studied from the point of view of map coloring theorems (see for example, [16, 44]), where non-orientable surfaces are also considered. Graphs on non-orientable surfaces play an important role in the study of moduli spaces of real algebraic curves [15]. In this section we use the notion of *Möbius graphs* following [34, 36] to emphasize that our graphs are on *unoriented* surfaces, which may or may not be orientable.

A ribbon graph is a graph with a cyclic order given at each vertex to the set of half-edges incident to it. Equivalently, it is a graph drawn on a closed oriented surface [32]. Vertices with cyclic orders are connected by ribbon like edges to form a ribbon graph [22]. In order to deal with a graph on a non-orientable surface, it is necessary to allow ribbons with a *twist*. We can use the group $\mathfrak{S}_2 = \{\pm 1\}$ to indicate a twist of an edge with the value -1 and no twist with 1 . We are thus led to consider a ribbon graph with $\mathfrak{S}_2$ -color at every edge. A *vertex flip* operation of an edge-colored ribbon graph is a move at a vertex V of a graph that reverses the cyclic order at V and interchanges the color of the edge that is incident to V . If an edge makes a loop and incident to V twice, then its color is changed twice and hence preserved. Two $\mathfrak{S}_2$ -colored ribbon graphs are *equivalent* if one is brought to the other by a sequence of vertex flip operations. A *Möbius graph* is the equivalence class of a $\mathfrak{S}_2$ -colored ribbon graph [36].

Let Γ_c be a connected $\mathfrak{S}_2$ -colored ribbon graph, and G a compact Lie group. For every vertex V of Γ_c , we label the half-edges incident to V as $(V1, V2, \dots, Vn)$ according to the cyclic order, where n is the valence (or degree) of the vertex. We call the expression

$$(6.1) \quad \Phi(V) = \delta(x_{V1} \cdots x_{Vn}) \in \mathcal{D}'(G^n)$$

the *vertex contribution*. It is invariant under the cyclic permutation of the half-edges. Suppose an edge E is incident to vertices V and W of Γ_c , and consists of two half-edges Vi and Wj . The *propagator* of the edge E is the expression

$$(6.2) \quad \Phi(E) = \delta(x_{Vi} x_{Wj}^c) \in \mathcal{D}'(G^2),$$

where $c \in \mathfrak{S}_2 = \{\pm 1\}$ is the color of E . A few remarks are necessary for the definition of a vertex contribution and a propagator as distributions. For every $n \geq 2$, the multiplication

$$m_n : G^n \ni (x_1, \dots, x_n) \longmapsto x_1 \cdots x_n \in G$$

is a real analytic map and

$$\psi_n : G^n \ni (x_1, \dots, x_{n-1}, x_n) \xrightarrow{\sim} ((x_1, \dots, x_{n-1}), (x_1 \cdots x_{n-1}) \cdot x_n) \in G^{n-1} \times G$$

is a real analytic isomorphism. The Jacobian $J(x_1, \dots, x_{n-1}, y)$ of the isomorphism ψ_n gives the push-forward measure

$$(\psi_n)_*(dx_1 \cdots dx_n) = J(x_1, \dots, x_{n-1}, y) dx_1 \cdots dx_{n-1} dy .$$

The vertex contribution (6.1) is a distribution defined by the continuous linear functional

$$C^\infty(G^n) \ni f(x_1, \dots, x_n) \longmapsto \int_{G^{n-1}} J(x_1, \dots, x_{n-1}, 1) f(x_1, \dots, x_{n-1}, (x_1 \cdots x_{n-1})^{-1}) dx_1 \cdots dx_{n-1} \in \mathbb{C} .$$

The propagator as a distribution is defined similarly. We note that the product of all vertex contributions and the product of all propagators

$$\prod_{V \in \Gamma_c} \Phi(V) \quad \text{and} \quad \prod_{E \in \Gamma_c} \Phi(E)$$

are well-defined distributions in $\mathcal{D}'(G^{2e})$, where $e = e(\Gamma_c)$ is the number of edges of the graph.

Now we come to the heuristic part of the argument. Let us “define”

$$G_{\Gamma_c} = \int_{G^{2e(\Gamma_c)}} \prod_{V \in \Gamma_c} \Phi(V) \prod_{E \in \Gamma_c} \Phi(E) dx ,$$

where dx is the product measure on $G^{2e(\Gamma_c)}$. Although these products are individually well-defined distributions on the space $G^{2e(\Gamma_c)}$, their product does not make sense in general. For now, let us suppose that G_{Γ_c} is legally defined. Then the first thing we can show is that *it is invariant under vertex flip operations*. To see this, choose a vertex V at which we perform the vertex flip. Suppose it is incident to half-edges $V1, \dots, Vn$ in this cyclic order. We denote by $V'i$ the other half-edge connected to Vi . Of course we do not mean they are all incident to another vertex labeled by V' . This is just a notation for the other half. Then the vertex flip invariance follows from

$$\begin{aligned} \delta(x_{V'1}^{-c} \cdots x_{V'n}^{-c}) &= \int_{G^n} \delta(x_{V1} \cdots x_{Vn}) \delta(x_{V1} x_{V'1}^c) \cdots \delta(x_{Vn} x_{V'n}^c) dx_{V1} \cdots dx_{Vn} \\ &= \int_{G^n} \delta(x_{Vn} \cdots x_{V1}) \delta(x_{V1} x_{V'1}^{-c}) \cdots \delta(x_{Vn} x_{V'n}^{-c}) dx_{V1} \cdots dx_{Vn} \\ &= \delta(x_{V'n}^c \cdots x_{V'1}^c) , \end{aligned}$$

where the first and the last delta functions are the same because of (5.8). Notice that the vertex flip invariance of G_{Γ_c} implies that this quantity is associated to a Möbius graph. So we define

$$(6.3) \quad G_\Gamma = \int_{G^{2e(\Gamma)}} \prod_{V \in \Gamma} \Phi(V) \prod_{E \in \Gamma} \Phi(E) dx$$

for every Möbius graph Γ .

The second property we can show, still assuming (6.3) being well-defined, is that *it is invariant under an edge contraction*. As explained in [32, 36], an edge contraction of a graph removes an edge that is incident to two distinct vertices and put these vertices together. If the edge is not twisted, then this is exactly the same edge contraction of [32]. If the edge is

twisted, then we first apply a vertex flip operation to one of the vertices to untwist the edge, and then proceed with the usual edge contraction. To see the invariance, suppose E is the edge to be removed, which is incident to two vertices $V \neq W$. Without loss of generality, we assume that E is not twisted, as mentioned above. Let us denote by $\delta(xy)$ the propagator. Using cyclic permutations of vertex contributions, we can assume that $\delta(vx)$ and $\delta(yw)$ are the vertex contributions at V and W , respectively, where v and w are products of group elements. The invariance under the edge contraction then follows from

$$(6.4) \quad \int_{G^2} \delta(vx)\delta(xy)\delta(yw)dx dy = \int_G \delta(vy^{-1})\delta(yw)dy = \delta(vw) .$$

Note that $\delta(vw)$ is the vertex contribution of the new vertex obtained by joining V and W together.

Recall the fact that every connected Möbius graph Γ determines a unique closed surface S_Γ and its cell-decomposition. Let $v(\Gamma)$, $e(\Gamma)$, and $f(\Gamma)$ denote the number of 0, 1, and 2-cells, respectively. The Euler characteristic of the surface is given by

$$\chi(S_\Gamma) = v(\Gamma) - e(\Gamma) + f(\Gamma) .$$

If Γ is orientable, then S_Γ is an orientable surface of genus g such that $\chi(S_\Gamma) = 2 - 2g$, and if Γ is non-orientable, then S_Γ is a non-orientable surface of cross-cap genus k with $\chi(S_\Gamma) = 2 - k$. Notice that the edge contraction preserves the Euler characteristic and the number $f(\Gamma)$ of 2-cells. Using the same argument of [36], we can calculate G_Γ for every graph. If the graph is orientable of genus $g \geq 1$, then

$$(6.5) \quad G_\Gamma = |G|^{f(\Gamma)-1} \int_{G^{2g}} \delta([x_1, y_1] \cdots [x_g, y_g]) dx_1 dy_1 \cdots dx_g dy_g ,$$

and if it is non-orientable of cross-cap genus k , then

$$(6.6) \quad G_\Gamma = |G|^{f(\Gamma)-1} \int_{G^k} \delta(x_1^2 \cdots x_k^2) dx_1 \cdots dx_k .$$

Regardless the orientation of the graph, we have a general formula

$$(6.7) \quad G_\Gamma = |G|^{f(\Gamma)-1} |\text{Hom}(\pi_1(S_\Gamma), G)|$$

if the expressions (6.5) and (6.6) are well-defined. In the sense of (6.7), we have the alternative “definition” (6.3) of the volume of the representation variety.

It is clear from the invariance under edge contraction that G_Γ of (6.3) is always ill-defined if $\chi(S_\Gamma) = 2$. As we have seen in Section 4, the integral (6.6) is ill-defined in general for graphs with $\chi(S_\Gamma) = 1$. For an orientable graph of genus $g \geq 2$, the integral (6.5) makes sense only when the group G is semisimple (see Section 3).

Everything we discussed in this section makes sense for a finite group G , and we have the Feynman diagram expansion formula

$$(6.8) \quad \log \int_{\{x \in \mathbb{R}[G] \mid x^* = x\}} e^{-\frac{1}{4}\langle x^2 \rangle} e^{\sum_j \frac{t_j}{2j} \langle x^j \rangle} d\mu(x) = \sum_{\substack{\Gamma \text{ connected} \\ \text{Möbius graph}}} \frac{1}{|\text{Aut}_M(\Gamma)|} G_\Gamma \prod_j t_j^{v_j(\Gamma)}$$

established in [35, 36], where $\langle \rangle = \frac{1}{|G|} \chi_{\text{reg}}$ is the normalized trace of the finite-dimensional von Neumann algebra $\mathbb{R}[G]$, which is the real group algebra of G , and $v_j(\Gamma)$ is the number of j -valent vertices of Γ . We note that if we use the regular representation itself instead of the normalized trace in the integral, then G_Γ of the graphical expansion is replaced by the familiar $|G|^{\chi(S_\Gamma)} |\text{Hom}(\pi_1(S_\Gamma), G)|$. We refer to the above mentioned articles for

more explanation of the expansion formula. The point we wish to make here is that the RHS of (6.8) *tends to be* well-defined for a compact semisimple Lie group G when the Euler characteristic $\chi(S_\Gamma)$ is negative. Then how should we consider the integral of the LHS for a compact Lie group? Is the ill-defined nature of the infinite dimensional integral concentrated on the non-hyperbolic surfaces? Namely, if we remove the contributions of the surfaces of non-negative Euler characteristics from the integral over the infinite-dimensional space, would then the integral become well-defined? If the integral over the real group algebra $L^2_\mathbb{R}(G)$ or even $\mathcal{D}'_\mathbb{R}(G)$ could make sense in some way, then Peter-Weyl theory gives a decomposition of the integral into an infinite sum of matrix integrals, and the volume formulas (3.7) and (4.7) should follow from the comparison of terms corresponding to orientable and non-orientable surfaces in the Feynman diagram expansion of the integral, as it was the case for a finite group [35, 36].

The alternative definition (6.3) suggests that this subject is related to the study of tensor categories and modular functors [2, 7].

REFERENCES

- [1] J. J. Andrews and M. L. Curtis, *Free groups and handlebodies*, Proceedings of the American Mathematical Society **16** (1965), 192–195.
- [2] Bojko Bakalov and Alexander Kirillov, Jr., *Lectures on tensor categories and modular functors*, University Lecture Series **21**, American Mathematical society, 2001.
- [3] G. V. Belyi, *On galois extensions of a maximal cyclotomic fields*, Math. U.S.S.R. Izvestija **14** (1980), 247–256.
- [4] D. Bessis, C. Itzykson and J. B. Zuber, *Quantum field theory techniques in graphical enumeration*, Advanced in Applied Mathematics **1** (1980), 109–157.
- [5] C. Brézin, C. Itzykson, G. Parisi, and J.-B. Zuber, *Planar diagrams*, Communications in Mathematical Physics **59** (1978), 35–51.
- [6] Theodor Bröcker and Tammo tom Dieck, *Representations of compact Lie groups*, Springer-Verlag, 1985.
- [7] Vyjayanthi Chari and Andrew Pressley, *A guide to quantum groups*, Cambridge University Press, 1995.
- [8] Marc Culler and Peter B. Shalen, *Varieties of group representations and splittings of 3-manifolds*, Annals of Mathematics **117** (1983), 109–146.
- [9] Richard P. Feynman, *Space-time approach to quantum electrodynamics*, Physical Review **76** (1949), 769–789.
- [10] Daniel S. Freed and Frank Quinn, *Chern-Simons theory with finite gauge group*, Communications in Mathematical Physics **156** (1993), 435–472.
- [11] Georg Frobenius, *Über Gruppencharaktere*, Sitzungsberichte der königlich preussischen Akademie der Wissenschaften (1896), 985–1021.
- [12] Georg Frobenius and Isaai Schur, *Über die reellen Darstellungen der endlichen Gruppen*, Sitzungsberichte der königlich preussischen Akademie der Wissenschaften (1906), 186–208.
- [13] I. M. Gelfand, M. I. Graev, G. E. Shilov, and N. Ya. Vilenkin, *Generalized functions*, volume 1–5, Academic Press, New York, 1964–1968.
- [14] William Goldman, *Ergodic theory on moduli spaces*, Annals of Mathematics **146** (1997), 475–507.
- [15] I. P. Goulden, J. L. Harer, J. L. and D. M. Jackson, *A geometric parametrization for the virtual Euler characteristics of the moduli spaces of real and complex algebraic curves*, Trans. Amer. Math. Soc. **353** (2001), 4405–4427.
- [16] Jonathan L. Gross and Thomas W. Tucker, *Topological graph theory*, John Wiley & Sons, 1987.
- [17] Alexander Grothendieck, *Esquisse d’un programme* (1984), reprinted in [46], 7–48.
- [18] John L. Harer, *The virtual cohomological dimension of the mapping class group of an orientable surface*, Inventiones Mathematicae **84** (1986), 157–176.
- [19] John L. Harer and Don Zagier, *The Euler characteristic of the moduli space of curves*, Inventiones Mathematicae **85** (1986), 457–485.
- [20] Allen Hatcher, *On triangulations of surfaces*, Topology and its Applications **40** (1991), 189–194.
- [21] Gareth A. Jones, *Characters and surfaces: a survey*, London Mathematical Society Lecture Note Series **249**, *The atlas of finite groups: ten years on*, Robert Curtis and Robert Wilson, Eds., (1998), 90–118.
- [22] Maxim Kontsevich, *Intersection theory on the moduli space of curves and the matrix Airy function*, Communications in Mathematical Physics **147** (1992), 1–23.
- [23] Peter B. Kronheimer and Tom S. Mrowka, *Embedded surfaces and the structure of Donaldson’s polynomial invariants*, Journal of Differential Geometry **41** (1995), 573–734.

- [24] Greg Kuperberg, *Involutory Hopf algebras and 3-manifold invariants*, International Journal of Mathematics **2** (1991), 41–66.
- [25] Kefeng Liu, *Heat kernel and moduli space*, Mathematical Research Letters **3** (1996), 743–762.
- [26] Kefeng Liu, *Heat kernel and moduli space II*, Mathematical Research Letters **4** (1996), 569–588.
- [27] Kefeng Liu, *Heat kernels, symplectic geometry, moduli spaces and finite groups*, Surveys in Differential Geometry **5** (1999), 527–542.
- [28] Saunders Mac Lane, *Categories for the working mathematician*, Springer-Verlag, 1971.
- [29] A. D. Mednykh, *Determination of the number of nonequivalent coverings over a compact Riemann surface*, Soviet Mathematics Doklady **19** (1978), 318–320.
- [30] Motohico Mulase, *Asymptotic analysis of a hermitian matrix integral*, International Journal of Mathematics **6** (1995), 881–892.
- [31] Motohico Mulase, *Lectures on the asymptotic expansion of a hermitian matrix integral*, in Supersymmetry and Integrable Models, Henrik Aratin et al., Editors, Springer Lecture Notes in Physics **502** (1998), 91–134.
- [32] Motohico Mulase and Michael Penkava, *Ribbon graphs, quadratic differentials on Riemann surfaces, and algebraic curves defined over $\mathbb{Q}$* , Asian Journal of Mathematics **2** (1998), 875–920.
- [33] Motohico Mulase and Michael Penkava, *Periods of Strebel differentials and algebraic curves defined over the field of algebraic numbers*, math.AG/0107119 (2001), to appear in Asian Journal of Mathematics.
- [34] Motohico Mulase and Andrew Waldron, *Duality of orthogonal and symplectic matrix integrals and quaternionic Feynman graphs*, math-ph/0206011 (2002).
- [35] Motohico Mulase and Josephine T. Yu, *A generating function of the number of homomorphisms from a surface group into a finite group*, math.QA/0209008 (2002).
- [36] Motohico Mulase and Josephine T. Yu, *Non-commutative matrix integrals and representation varieties of surface groups in a finite group*, math.QA/0211127 (2002).
- [37] M. S. Narasimhan and C. S. Seshadri, *Holomorphic vector bundles on a compact Riemann surface*, Mathematische Annalen **155** (1964), 69–80.
- [38] M. S. Narasimhan and C. S. Seshadri, *Stable and unitary vector bundles on a compact Riemann surface*, Annals of Mathematics **82** (1965), 540–567.
- [39] Andrei Okounkov, *Random matrices and random permutations*, math.CO/9903176 (1999).
- [40] Andrei Okounkov and Rahul Pandharipande, *Gromov-Witten theory, Hurwitz numbers, and matrix models*, I, math.AG/0101147 (2001).
- [41] Andrei Okounkov and Rahul Pandharipande, *The equivariant Gromov-Witten theory of $\mathbb{P}^1$* , math.AG/0207233 (2002).
- [42] Robert C. Penner, *Perturbation series and the moduli space of Riemann surfaces*, Journal of Differential Geometry **27** (1988), 35–53.
- [43] Doug Pickrell and Eugene Z. Xia, *Ergodicity of mapping class group actions on representation varieties, I. Closed surfaces*, math.DS/0006098 (2000).
- [44] Gerhard Ringel, *Map color theorem*, Springer-Verlag, 1974.
- [45] Leila Schneps, *The grothendieck theory of dessins d'enfants*, London Mathematical Society Lecture Notes Series, vol. 200, 1994.
- [46] Leila Schneps and Pierre Lochak, editors, *Geometric Galois actions: Around Grothendieck's esquisse d'un programme*, London Mathematical Society Lecture Notes Series, vol. 242, 1997.
- [47] Kurt Strebel, *Quadratic differentials*, Springer-Verlag, 1984.
- [48] Gerard 't Hooft, *A planar diagram theory for strong interactions*, Nuclear Physics B **72** (1974), 461–473.
- [49] V. S. Varadarajan, *Harmonic analysis on real reductive groups*, Lecture Notes in Mathematics **576**, Springer-Verlag, 1977.
- [50] V. S. Varadarajan, *Lie groups, Lie algebras, and their representations*, Springer-Verlag, 1984.
- [51] Garth Warner, *Harmonic analysis on semi-simple Lie groups I*, Springer-Verlag, 1972.
- [52] Edward Witten, *On quantum gauge theories in two dimensions*, Communications in Mathematical Physics **141** (1991), 153–209.
- [53] Edward Witten, *Two dimensional gravity and intersection theory on moduli space*, Surveys in Differential Geometry **1** (1991), 243–310.
- [54] Edward Witten, *Two dimensional gauge theory revisited*, Journal of Geometry and Physics **9** (1992), 303–368.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CA 95616–8633
E-mail address: mulase@math.ucdavis.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, EAU CLAIRE, WI 54702–4004
E-mail address: penkavmr@uwec.edu